

BEFORE THE PUBLIC UTILITIES COMMISSION  
OF THE STATE OF SOUTH DAKOTA

IN THE MATTER OF THE APPLICATION OF  
MONTANA-DAKOTA UTILITIES CO. AND  
OTTER TAIL POWER COMPANY FOR A  
PERMIT TO CONSTRUCT THE BIG STONE  
TO SOUTH ELLENDALE 345KV  
TRANSMISSION LINE

DIRECT TESTIMONY OF  
GREGORY TYLKA

EL13-028

**1. State your name and address.**

Gregory L. Tylka, 922 New Hampshire Circle, Ames, IA 50014

**2. Describe your occupation and educational background.**

I am a professor in the Department of Plant Pathology and Microbiology at Iowa State University. My responsibilities are to conduct research and extension educational programs on the biology and management of plant-parasitic nematodes, with a heavy emphasis on the soybean cyst nematode (SCN), *Heterodera glycines*.

**3. Is a complete copy of your resume or C.V. attached to this testimony?**

Yes.

**4. Are you familiar with the proposed B.S.S.E. transmission line?**

Yes.

**5. Could the construction and maintenance of the proposed line impact the spread of soil-born pests like the Soybean Cyst Nematode or "SCN"?**

Yes, construction and maintenance of the proposed line could accelerate the spread of SCN into previously uninfested townships, farms, and fields.

**6. Please explain what the SCN is.**

Nematodes are microscopic worms, many of which live in terrestrial ecosystems such as row-crop farm fields. A majority of soil-dwelling nematodes are not harmful to plants. They consume bacteria, fungi, and other nematodes and contribute to cycling of nutrients in the soil. But agricultural soils also contain nematodes called plant-parasitic nematodes that feed on plants. Many plant-parasitic nematodes are believed to be native to midwestern soils, feeding off of naturally occurring plants in the landscape. These native, plant-parasitic nematodes must reach high population densities (numbers) in order to be damaging to crops. The soybean cyst nematode, SCN, is very different than native plant-parasitic nematodes that reside in typical midwestern soils. SCN is an introduced pest, which means it

does not naturally occur in the United States. Introduced pests like SCN have few or no natural enemies when they are first introduced into a new environment, and the lack of natural enemies allows for very high reproduction in the first few years following introduction into an area. SCN is believed to have been introduced into North America from Asia (Riggs, 2004). The nematode feeds from the soybean root's vascular tissue, stunting the roots and making the roots less able to use atmospheric nitrogen, making it highly damaging to the soybean crop. Also, soybean plants infected with SCN are more vulnerable and suffer greater yield loss from soil-borne soybean diseases caused by fungi that are commonly found in soybean fields in the Midwest (Niblack et al., 2006). The nematode has a relatively short generation time (24 to 40 days), allowing it to complete three to six generations on a single soybean crop, and the nematode females each produce 250 or more eggs, allowing SCN population densities to increase very quickly. Finally, eggs of SCN can survive in a dormant condition in the soil without soybeans or another host crop being grown for a decade or more (Riggs, 2004).

**7. Please describe the research you've done, or of which you are aware, regarding the impact and spread of SCN.**

The soybean cyst nematode, scientific name *Heterodera glycines*, is considered the most damaging pathogen of soybean in the United States and Canada. Annual yield losses in the United States alone are estimated to exceed \$1 billion (Wrather et al., 2010). Soybean cyst nematode can reduce soybean yields without causing aboveground symptoms (Wang et al., 2003), which allows the nematode to build up in fields to more damaging levels before farmers notice a problem. In a single growing season, SCN population densities can increase from less than 1,500 eggs per 100 cm<sup>3</sup> of soil (a little less than a half cup) in the spring to more than 19,000 eggs per 100 cm<sup>3</sup> of soil on susceptible soybean varieties and nearly 6,000 eggs per 100 cm<sup>3</sup> of soil on SCN-resistant soybean varieties at the time of harvest (Tylka et al., 2013). The map of the known distribution of SCN in North America was recently updated, and South Dakota was identified as a state that was particularly vulnerable to having SCN spread considerably in the future (Tylka and Marett, 2014).

**8. Please describe the impact SCN can have on crop production.**

It is difficult to scientifically quantify the exact impact of SCN on soybean yields in infested fields because yield loss is dependent on the nematode population density (with greater yield loss occurring at higher SCN numbers), the soybean variety grown (SCN-susceptible varieties suffer greater damage than SCN-resistant varieties), and weather (there is much greater SCN reproduction and greater yield loss in hot dry years than in years of moderate temperatures and adequate to excess moisture). However, one can gain some insight into the yield-reducing effects of SCN by comparing the yields of SCN-resistant and SCN-susceptible soybean varieties in experiments conducted in SCN-infested fields. SCN-resistant soybean varieties allow some nematode reproduction and, consequently the varieties suffer some yield loss. But SCN-resistant soybean varieties suppress reproduction of the nematode compared to susceptible soybeans, which allow unchecked nematode reproduction. My research program at Iowa State University compares the yields of >60 SCN-resistant soybean varieties to yields of several widely grown SCN-susceptible soybean varieties each year at nine locations throughout Iowa (see [www.isuscntrials.info](http://www.isuscntrials.info)). In 2013, the average yield of SCN-resistant soybeans ranged from 3.7 bushels per acre (5.6%) to 18.4 bushels per acre (56%) greater than the susceptible soybean varieties among the nine experiments (Tylka, 2014). Those yield differences equate to \$54 to \$268 per acre at current soybean prices (\$14.60 per bushel). At very high SCN population densities and under very stressful weather conditions (severe drought), yield loss can approach or exceed 50%.

**9. Please describe the ways that SCN can spread from field-to-field.**

The life stage of SCN that best survives dormant in the soil and poses the greatest threat for spread is the cyst, which is the egg-filled body of a dead SCN female. SCN females and cysts are about the size of a period at the end of a printed sentence in a newspaper, and each female and cyst can contain 200 or more eggs. Anything that moves soil particles of this size is capable of moving SCN. Common avenues of spreading the nematode include moving soil on farming equipment, wind-blown soil, and soil moved with surface erosion due to rainfall. Equipment that digs into and disrupts the soil to a depth of a foot or more would likely be more effective at spreading SCN than surface soil adhering to the wheels of vehicles simply because of the volume of soil being moved.

**10. Could both construction equipment and ordinary farm equipment cause SCN to spread?**

Yes. Movement of any equipment (example: tractors, sprayers, combines, cranes, cement trucks) could directly move SCN by transporting clumps of soil containing SCN cysts, which are the size of a period on a printed page and can be full of hundreds of eggs each.

**11. Has SCN been identified in the areas of South Dakota where the proposed B.S.S.E. Line would be constructed?**

Yes. SCN has been found in northeastern South Dakota as well as in southeastern North Dakota (Tylka and Marett, 2014). Also, because fields infested with SCN may not exhibit obvious symptoms of damage for years (Wang et al., 2003), it is likely that more fields and counties are infested with SCN than officially reported.

When SCN becomes first established in a new field, it tends to be aggregated or clustered in discrete areas in the field because the nematode is relatively immobile. The SCN second-stage juveniles that hatch from the egg are the only mobile and infective stage of the nematode, and these juveniles can move no more than an inch or so under their own power. So SCN usually takes years to spread throughout a field. But SCN reproduction rates (increases in numbers over time) on soybeans usually are greatest in the first few years after the nematode is introduced into a field because there are no natural enemies present in the soil (since SCN has never occurred in that field before) and the nematode will have good nutrition because the soybean crop will be relatively healthy.

**12. Can construction equipment used in a project like the proposed B.S.S.E. Line cause SCN to spread farther or more rapidly than ordinary farming practices? If so, how?**

Yes. Soil disturbed by construction equipment would likely result in greater spread of the nematode than soil disturbed by most other common occurrences by making the soil more friable (easily crumbled) and prone to erosion compared to soil that is left undisturbed or disturbed just minimally.

Also, soil moved by construction equipment could be from properties owned by various farmers. And SCN-infested soil from a less-than-diligent farmer's field could be moved into a field farmed by someone who has diligently worked to avoid introduction of SCN into their fields by careful management of the movement of soil.

**13. Could ongoing maintenance of the proposed B.S.S.E. line also impact the spread of SCN in the region? If so, how?**

Ongoing maintenance of the proposed line would not likely have much greater impact on spreading SCN than other activities involving vehicles traveling through the SCN-infested fields, but it depends somewhat on the condition of the soil at the time that vehicles or crews are present in the field. The least chance of movement, in my opinion, would be when the soil is frozen because frozen soil would not adhere well to vehicles or on people's boots. And the greatest chance would be when the soil is moist enough to allow for mud to easily adhere to vehicles and on people's boots.

**14. Are there ways to remove SCN from a field once it has been introduced? If so, please describe these.**

Once SCN is introduced into a field, there is nothing that can be done to eradicate the nematode other than to not grow soybeans for an extended period of time. Many of the eggs that are present within cysts (dead females) of SCN are in a dormant state and capable of surviving in the absence of a host crop for a decade or more. There are anecdotal reports of SCN surviving in soil without a host for thirty years (Riggs, 2004). So a field that is infested with SCN would have to remain fallow or be planted with a nonhost crop from 10 to 30 years or more to eliminate the nematode from the field.

**15. Are there ways to mitigate the damage caused by SCN in a field once it has been introduced? If so, please describe these.**

Soil applied chemicals to kill SCN directly are no longer available for use in fields in the Midwest. Current management options are 1) minimize field-to-field movement of SCN-infested soil, 2) grow SCN-resistant soybean varieties, 3) grow nonhost crops, and 4) use seed-applied nematode protectants when planting soybeans.

The most effective option to maximize soybean production is delaying introduction of the nematode into a field or an area. States like South Dakota and North Dakota are in a unique position to be able to significantly delay the spread of SCN into the soybean-producing areas of their states by managing the movement of SCN-infested soil from field to field and farm to farm.

Growing nonhost crops, such as corn, in an SCN-infested field will reduce SCN numbers, but the reduction can vary from 5 or 10% to about 50% within a single growing season. Unfortunately, the reduction in egg numbers as a consequence of growing a nonhost crop does not occur in multiple years in a row. That is, the greatest reduction of SCN egg numbers occurs in the first year that a nonhost crop is grown, with a slight reduction in numbers occurring the second year, and very little reduction occurring in years thereafter because dormant eggs will primarily remain in the soil after two years of growing a nonhost crop.

SCN-resistant soybean varieties can be effective at producing acceptable soybean yields in SCN-infested fields and slowing the build-up of the nematode, but as described above, even SCN-resistant soybean varieties suffer some yield loss. Also, there will be considerably fewer SCN-resistant soybean varieties adapted for growing in South Dakota than in more southern areas of the Midwest, including Iowa. Another significant shortcoming of SCN-resistant soybean varieties is that almost all (>95%)

contain the same set of resistance genes (Tylka and Mullaney, 2013). This lack of genetic diversity has led to SCN populations in the Midwest developing an increased ability to reproduce on the SCN-resistant soybean varieties.

There are at least four nematode-protectant seed treatments that are being advertised as providing protection against early season infection by SCN. But the seed treatments infrequently increased yields or reduced SCN population densities in university field experiments conducted throughout the Midwest in 2012 and 2013. So the utility of these nematode-protectant seed treatments has yet to be proven.

**16. Are there ways to prevent SCN from being spread from field-to-field by construction or farm equipment? If so, please describe these.**

Fields could be tested for SCN in advance of moving equipment in by collecting soil samples from the fields and having the samples tested for presence of the nematode. But when we did follow-up testing of soil samples that tested negative for SCN that were submitted to the Iowa State University Plant and Insect Diagnostic Clinic, we discovered a 14% rate of false negative results (Tylka and Flynn, 2000). That is, 14% of the time, we observed SCN females growing on soybean roots after 30 days growing in leftover soil from samples that had tested negative for SCN with our standard extraction procedure. This rate of false negative results would likely occur with any laboratory processing the samples and is the result of soil clods containing SCN cysts remaining clumped during the soil processing procedure and not releasing the SCN cysts to be trapped on the sieves used in the process.

Another possible way to reduce likelihood of spread of SCN on equipment is to clean the equipment before it moves from field to field. Soil adhering to all parts of all machines must be washed off (although disinfecting probably isn't warranted). But this washing effort, no matter how thorough, can be inefficient because all of the soil on a vehicle must be removed and then the run-off water and soil from the rinsing must be directed away from the next area or field that will be worked in.

**17. Is there anything else you feel the Commission should know about SCN as it relates to the construction of the proposed B.S.S.E. transmission line?**

Following are references to scientific articles and extension publications that support specific statements made above:

Niblack, T.L., K.N. Lambert, and G.L. Tylka. 2006. A model plant pathogen from the kingdom Animalia: *Heterodera glycines*, the soybean cyst nematode. Annual Review of Phytopathology 44:283-303.

Riggs, R.D. 2004. History and distribution. Pages 9-39 in: Biology and Management of Soybean Cyst Nematode: Second Edition. Walsworth Publishing Company, Marceline, MO.

Tylka, G.L. and P.H. Flynn. 2000. Effectiveness of soil analysis for presence of the soybean cyst nematode, *Heterodera glycines*. Journal of Nematology 32: 467-468.

Tylka, G.L. and M.P. Mullaney. 2013. Soybean cyst nematode-resistant soybeans for Iowa. Iowa State University Extension Publication PM 1649, 22 pp.

Tylka, G.L., G.D. Gebhart, C.C. Marett, and M.P. Mullaney. 2013. Evaluation of soybean varieties resistant to soybean cyst nematode in Iowa – 2012. Iowa State University Extension, publication IPM-52, 32 pp.

Tylka, G.L. and C.C. Marett. 2014. Distribution of the soybean cyst nematode (*Heterodera glycines*) in the United States and Canada: 1954 to 2014. Plant Health Progress (accepted for publication).

Tylka, G. 2014. Trial results show dual benefits of SCN resistance. Iowa State University Integrated Crop Management News, January 17, 2014 (<http://www.extension.iastate.edu/CropNews/2014/00117tylka.htm>).


Wang, J., T.L. Niblack, J.N. Tremaine, W.J. Wiebold, G.L. Tylka, C.C. Marett, G.R. Noel, O. Myers, and M.E. Schmidt. 2003. The soybean cyst nematode reduces soybean yield without causing obvious symptoms. Plant Dis. 87:623-628.

Wrather, A., G. Shannon, R. Balardin, L. Carregal, R. Escobar, G.K. Gupta, Z. Ma, W. Morel, D. Ploper, and A. Tenuta. 2010. Effect of diseases on soybean yield in the top eight producing countries in 2006. Online. Plant Health Progress doi:10.1094/PHP-2010-0102-01-RS.

**18. Does this conclude your testimony?**

Yes.

Dated this 23<sup>rd</sup> day of April, 2014


  
Gregory Tylka

STATE OF Iowa )  
                                  :SS  
COUNTY OF Story )

On this 23 day of April, 2014, before me personally appeared Gregory Tylka, known to me to be the person who is described in, and who executed the foregoing instrument and acknowledged to me that he or she executed the same.

(seal)



  
Rita Brucland, Notary Public  
My Commission Expires: Nov. 14, 2015

**CERTIFICATE OF SERVICE**

The undersigned attorney hereby certifies that at true and correct copy of the foregoing **DIRECT TESTIMONY OF GERALD PESALL** was served upon the following parties of record,

electronically or in paper form, this 24<sup>th</sup> day of April, 2014:

Thomas J. Welk, Attorney  
P.O. Box 5015  
Sioux Falls, SD 57117-5015  
[tjwelk@bgpw.com](mailto:tjwelk@bgpw.com)

Patricia Van Gerpen  
Executive Director  
South Dakota P.U.C.  
500 E. Capitol Ave.  
Pierre, SD 57501  
[patty.vangerpen@state.sd.us](mailto:patty.vangerpen@state.sd.us)

Karen Cremer, Attorney  
500 E. Capitol Ave.  
Pierre, SD 57501  
[karen.cremer@state.sd.us](mailto:karen.cremer@state.sd.us)

Brian Rounds  
Staff Analyst  
South Dakota P.U.C.  
500 E. Capitol Ave.  
Pierre, SD 57501  
[brian.rounds@state.sd.us](mailto:brian.rounds@state.sd.us)

Darren Kearny  
Staff Analyst  
South Dakota P.U.C.  
500 E. Capitol Ave.  
Pierre, SD 57501  
[darren.kearny@state.sd.us](mailto:darren.kearny@state.sd.us)


Jennifer Smestad, Attorney  
214 S. Cascade St.  
Fergus Falls, MN 56538  
[jsmestad@ottertail.com](mailto:jsmestad@ottertail.com)

Daniel S. Kuntz, Attorney  
P.O. Box 5650  
1200 West Century Ave.  
Bismarck, ND 58506  
[dan.kuntz@mduresources.com](mailto:dan.kuntz@mduresources.com)

Maxine Fischer  
Brown County Auditor  
25 Market St., Ste. 1  
Aberdeen, SD 57401  
[maxine.fischer@browncounty.s  
d.gov](mailto:maxine.fischer@browncounty.sd.gov)

Sandra Raap  
Day County Auditor  
711 W. First St., Ste. 204  
Webster, SD 57274  
[dcaud@itctel.com](mailto:dcaud@itctel.com)

Karen Layher  
Grant County Auditor  
210 E. Fifth Ave.  
Milbank, SD 57252  
[karen.layher@state.sd.us](mailto:karen.layher@state.sd.us)

  
N. Bob Pesall, Attorney

## GREGORY L. TYLKA

Business Address:  
Department of Plant Pathology and Microbiology  
Iowa State University  
Ames, Iowa 50011  
(515) 294-3021

Home Address:  
922 New Hampshire Circle  
Ames, Iowa 50014  
(515) 509-3719

### EDUCATION

Ph.D. Degree, University of Georgia, Athens, Georgia. 1990, Summa Cum Laude. Major: Plant Pathology. Major professor: Dr. Richard S. Hussey. Dissertation title: The interactions of vesicular-arbuscular mycorrhizal fungi with *Heterodera glycines* and soil actinomycetes on soybean.

M.S. Degree, California University of Pennsylvania, California, Pennsylvania. 1985, Summa Cum Laude. Major: Biology. Major Professor: Dr. Barry B. Hunter. Thesis title: The isolation, quantification, and possible integrated control of *Cylindrocladium* species from Mont Alto and Penn forest tree nursery soils.

B.S. Degree, California University of Pennsylvania, California, Pennsylvania. 1983, Magna Cum Laude. Advisor: Dr. Samuel K. Hood. Major: Biology.

### EMPLOYMENT RECORD

July 2000 to present: Professor, Department of Plant Pathology and Microbiology, Iowa State University, Ames, Iowa. Responsibilities include graduate teaching and extension efforts concerning all plant-parasitic nematode species and research on effects of cultural practices and soybean resistance on soybean cyst nematode, *Heterodera glycines*, population densities and soybean yield suppression and the interactions of soybean insect and weed pests with soybean cyst nematode.

July 2003 to September 2010: Coordinator (half time), Corn and Soybean Initiative, College of Agriculture and Life Sciences, Iowa State University, Ames, Iowa. Responsibilities include providing planning, organization, and leadership to coordinate applied research on corn and soybean production in Iowa and the transfer of science-based crop production information to Iowa corn and soybean growers.

July 1995 to 2000: Associate Professor, Department of Plant Pathology, Iowa State University, Ames, Iowa. Responsibilities include extension efforts concerning all plant-parasitic nematode species and research on effects of cultural practices and soybean resistance and tolerance on soybean cyst nematode, *Heterodera glycines*, population densities and soybean yield suppression. Also investigating stimulation and inhibition of hatching of soybean cyst nematode eggs and the interactions of soybean insect and weed pests with soybean cyst nematode.

February 1990 to June 1995: Assistant Professor, Department of Plant Pathology, Iowa State University, Ames, Iowa. Primary responsibilities included research on the effects of cultural practices and soybean resistance and tolerance on soybean cyst nematode, *Heterodera glycines*,



### EMPLOYMENT RECORD (continued)

and soybean yield suppression. Also investigated induced hatching of soybean cyst nematode eggs and the influence of soybean insect and weed pests on soybean cyst nematode. Additionally, considerable extension and teaching responsibilities were components of this position.

August 1988 to December 1989: Half-time laboratory technician for Dr. Richard S. Hussey, Department of Plant Pathology, University of Georgia, Athens, Georgia. Participated in a research program utilizing monoclonal antibodies in immunocytochemical studies of the nature and function of disease-inducing secretions of the root-knot nematode, *Meloidogyne incognita*. Investigations were being conducted on the molecular basis of pathogenesis by identifying and characterizing the esophageal gland secretions within juvenile and adult nematodes that modify recipient host root tissue during nematode parasitism.

September 1985 to December 1989: Graduate Research Assistant, Department of Plant Pathology, University of Georgia, Athens, Georgia. Conducted field, greenhouse and laboratory investigations on the influence of vesicular-arbuscular mycorrhizal fungi (*Gigaspora* and *Glomus*) upon development and reproduction of the soybean cyst nematode, *Heterodera glycines*, on soybean. Also investigated effect of soil inhabiting *Streptomyces* species on mycorrhizal fungal spore germination and penetration and colonization of soybean roots.

September 1983 to August 1985: Graduate Research Assistant, Department of Biological and Environmental Sciences, California University of Pennsylvania, California, Pennsylvania. This project, partially funded by the Pennsylvania Bureau of Forestry, involved soil microbiological research into the biology of *Cylindrocladium* species and their infection of conifer seedling roots in Pennsylvania's State Forest Tree Nurseries. Served as teaching assistant for undergraduate Principles of Biology and Scientific Photography courses and instructor of Microscopy and Photography courses for the Summer Academy for Gifted Students (1984 and 1985).

### PROFESSIONAL AND HONORARY ASSOCIATIONS

American Phytopathological Society  
American Soybean Association  
Beta Beta Beta  
Iowa Corn Growers Association

Iowa Soybean Association  
Sigma Xi  
Society of Nematologists

### AWARDS

Regents Faculty Excellence Award, Board of Regents, State of Iowa (2010)

Educational Materials Award of Excellence for Computer Software Programs, American Society of Agronomy, for computer training module for certified crop advisors on biology and management of corn nematodes (2009)

AWARDS (continued)

- Educational Materials Award of Excellence for Printed Materials, American Society of Agronomy, for special issue of the Integrated Crop Management Newsletter (2007)
- Educational Materials Award of Excellence for Computer Software Programs, American Society of Agronomy, for computer training module for certified crop advisors on biology and management of the soybean cyst nematode (2006)
- Dean's Citation for Extraordinary Contributions to the College of Agriculture (2005)
- Outstanding Individual Achievement Award, Iowa State University Extension (2002)
- Meritorious Service Extension Education Award, United Soybean Board (2000)
- Excellence in Extension Award, American Phytopathological Society (1999)
- Meritorious Service Award, Iowa State University Extension (1999)
- Novartis Crop Protection Award, Society of Nematologists (1999)
- Meritorious Service Production Research Award, United Soybean Board (1998)
- Outstanding Alumnus Award, Beta Beta Beta, California University of Pennsylvania (1992)

GRANTS (last 5 years, 2009-present)

- Toward increased efficacy of soybean cyst nematode management tools. **G.L. Tylka** and S. Pandey. Funded by the Center for Arthropod Management Technologies for \$120,00 total over two years (2014-2015).
- Development of multiple pest resistant soybeans for breeding and research purposes using field and molecular tools. A.K. Singh, M. O'Neal, **G. Tylka**, G. MacIntosh, and A. Singh. Funded by the Iowa Soybean Association for \$310,348 over three years (2013-2016).
- Enhancing soybean yield through strategic use of soybean seed treatments for seedling disease and insect pest management. A. Robertson, G. Munkvold, **G. Tylka**, and E. Hodgson. Funded by the Iowa Soybean Association for \$409,456 over three years (2013-2016).
- Exploring soybean aphid and soybean cyst nematode interactions for improved integrated management in Iowa. M.E. O'Neal, **G.L. Tylka**, G. MacIntosh, E.W. Hodgson, and M. McCarville. Funded by the Iowa Soybean Association for \$339,417 over three years (2013-2016).
- Developing an integrated management and communication plan for sudden death syndrome. D. Mueller, L. Leandro, C. Bradley, M. Chilvers, **G. Tylka**, K. Wise, S. Cianzio, J. Faghihi, A. Tenuta, V. Ferris, D. Malvick, A. Fakhoury, and G. Hartman. Funded by the North Central Soybean Research Program for \$500,000 over three years (2013 – 2015).

## GRANTS (continued)

Modifying *Bradyrhizobium japonicum* to enhance nodulated soybean disease resistance. R. Peters, L. Leandro, A. Robertson, and **G. Tylka**. Funded by the Iowa Soybean Association for \$246,266 over two years (2013-2015).

Continuation of assessment of nematode control and yield of SCN-resistant soybean varieties in response to different soybean cyst nematode populations (HG Types). **G. Tylka**. Funded by the Iowa Soybean Association for \$444,129 over three years (2012–2015).

Determining soybean pest and pesticide interactions as a means to optimize soybean yield, E. Hodgson, A. Gassmann, and **G. Tylka**. Funded by the Iowa Soybean Association for \$201,206 over three years (2012–2015).

Characterization of the mechanisms involved in the SDS-SCN interaction to develop soybean lines with resistance to SDS and to SCN, L. Leandro, **G. Tylka**, S. Cianzio, and O. Radwan, S.R. Cianzio. Funded by the Iowa Soybean Association for \$127,251 over two years (2012–2014).

Exploring soybean aphid and soybean cyst nematode interactions for improved integrated management in Iowa. M.E. O’Neal, **G.L. Tylka**, G. MacIntosh, E.W. Hodgson, and M. McCarville. Funded by the Iowa Soybean Association for \$339,417 over three years (2011-2014).

Assessing nematode control and yield of SCN-resistant soybean varieties in response to different soybean cyst nematode populations (HG types), **G. Tylka**. Funded by the Iowa Soybean Association for \$613,720 for three years (2010-2012).

Determining the impact of multiple pests on soybean yield and grain composition, G. MacIntosh, M. O’Neal, **G. Tylka**, P. Pedersen, and F. Avendano. Funded by the Iowa Soybean Association for \$300,326 for three years (2007-2010).

Increasing Iowa soybean profitability by renewing interest in managing the soybean cyst nematode, **G.L. Tylka**. Funded by the Iowa Soybean Association for \$517,199 for four years (2006-2010).

Improving soybean profitability in Iowa by reducing the hidden effects of brown stem rot and its interaction with the soybean cyst nematode, G. Tabor and **G.L. Tylka**. Funded by the Iowa Soybean Association for \$216,760 over three years (2006 to 2009).

## INDUSTRY RESEARCH CONTRACTS

AMVAC  
BASF Plant Sciences  
Bayer Crop Sciences  
Bushvale Seeds

CENEX/Land O’Lakes  
Chemtura  
DeKalb Genetics Corporation  
DM Crop Research, Inc.

## INDUSTRY RESEARCH CONTRACTS (continued)

Divergence Company	Novartis Crop Protection, Inc.
DuPont Crop Protection	Pioneer Hi-Bred International, Inc.
Evolutionary Genomics Inc.	Stine Seed Company
JGL, Inc.	Stoller Enterprises Inc
LiphaTech, Inc.	Syngenta Crop Protection
MBS Genetics	TJ Technologies, Inc.
Monsanto	Valent

## AGENCIES AND ORGANIZATIONS ADVISED

BASF	LiphaTech, Inc.
Bayer CropScience	Monsanto
DNA Plant Technologies	North Central Soybean Research Program
Garst Seeds	Novartis Crop Protection, Inc.
Illinois Department of Agriculture	Pioneer Hi-Bred International, Inc.
Iowa Department of Agriculture	Syngenta
Iowa Soybean Association	United Soybean Board

## PUBLICATIONS

## I. Refereed research articles:

**Tylka, G.L.** and C.C. Marett. Distribution of the soybean cyst nematode (*Heterodera glycines*) in the United States and Canada: 1954 to 2014. Plant Health Progress (accepted).

McCarville, M.T., D.H. Soh, **G.L. Tylka**, and M.E. O'Neal. 2014. Aboveground feeding by soybean aphid, *Aphis glycines*, affects soybean cyst nematode, *Heterodera glycines*, reproduction belowground. PLoS ONE 9(1): e86415. doi:10.1371/journal.pone.0086415.

McCarville, M.T., C. Kanobe, M. O'Neal, G. MacIntosh and **G.L. Tylka**. 2012. Measuring the yield and fatty-acid response of soybean cultivars with seed oil low in linolenic acid to multiple biotic stresses. Crop Protection 42:210-216.

McCarville, M.T., M. O'Neal, **G.L. Tylka**, C. Kanobe and G. MacIntosh. 2012. A nematode, fungus, and aphid interact via a shared host plant: implications for soybean management. Entomologia Experimentalis et Applicata 143:55-66, doi: 10.1111/j.1570-7458.2012.01227.x.

**Tylka, G.L.**, A.J. Sisson, L.C. Jesse, J. Kennicker and C.C. Marett. 2011. Testing for plant-parasitic nematodes that feed on corn in Iowa 2000-2010. Online. Plant Health Progress doi:10.1094/PHP-2011-1205-01-RS.

## PUBLICATIONS (continued)

- Tylka, G.L.**, T.C. Todd, T.L. Niblack, A.E. MacGuidwin, and T. Jackson. 2011. Sampling for plant-parasitic nematodes in corn strip trials comparing nematode management products. *Plant Health Progress* doi:10.1094/PHP-2011-0901-01-DG.
- Rotundo, J.L., **G.L. Tylka**, and P. Pedersen. 2010. Source of resistance affects soybean yield, yield components, and biomass accumulation in *Heterodera glycines*-infested fields. *Crop Science* 50:2565–2574.
- Pedersen, P., **G.L. Tylka**, A.P. Mallarino, A.E. MacGuidwin, N.C. Koval, and C.R. Grau. 2010. Correlation between soil pH, *Heterodera glycines* population densities, and soybean yield. *Crop Science* 50:1458–1464, doi: 10.2135/cropsci2009.08.0432.
- Niblack, T.L., **G.L. Tylka**, P. Arelli, J. Bond, B. Diers, P. Donald, J. Faghihi, V.R. Ferris, K. Gallo, R.D. Heinz, H. Lopez-Nicora, R. Von Qualen, T. Welacky, and J. Wilcox. 2009. A standard greenhouse method for assessing soybean cyst nematode resistance in soybean: SCE08 (Standardized Cyst Evaluation 2008). Online. *Plant Health Progress* doi:10.1094/PHP-2009-0513-01-RV.
- Rogovsa, N., F.W. Blackmer, and **G.L. Tylka**. 2009. Soybean yield and soybean cyst nematode densities related to soil pH, soil carbonate concentrations, and alkalinity stress index. *Agronomy Journal* 101:1019-1026 (also online, doi:10.2134/agronj2008.0086x).
- Studham, M., G.C. MacIntosh, F. Avendaño, D. Soh, and **G.L. Tylka**. 2009. The soybean resistance gene *Rag1* does not protect against soybean cyst and root-knot nematodes. Online. *Plant Health Progress* doi:10.1094/PHP-2009-0401-01-BR.
- Charlson, D.V., K.R. Harkins, and **G.L. Tylka**. 2008. Relationship between juvenile hatching and acridine orange fluorescence of *Heterodera glycines* eggs. *Nematology* 10:603-610.
- Zasada, I.A., F. Avendano, Y. Li, T. Logan, H. Melakeberhan, S. Koenning, and **G.L. Tylka**. 2008. Potential of an alkaline-stabilized biosolid to manage nematodes: case studies on soybean cyst and root-knot nematodes. *Plant Disease* 92:4-13.
- Gavassoni, W.L., **G.L. Tylka**, and G.P. Munkvold. 2007. Relationships among tillage practices, dissemination, and spatial patterns of *Heterodera glycines* and soybean yield. *Plant Disease* 91:973-978.
- Tabor, G.M., **G.L. Tylka**, and C.R. Bronson. 2007. Genotypes A and B of *Cadophora gregata* differ in ability to colonize susceptible soybean. *Plant Disease* 91:574-580.
- Tabor, G.M., **G.L. Tylka**, and C.R. Bronson. 2006. Soybean stem colonization by genotypes A and B of *Cadophora gregata* increases with increasing population densities of *Heterodera glycines*. *Plant Disease* 90:1297-1301.

## PUBLICATIONS (continued)

- Donald, P.A., P.E. Pierson, S.K. St. Martin, P.R. Sellers, G.R. Noel, A.E. MacGuidwin, J. Faghihi, V.R. Ferris, C.R. Grau, D.J. Jardine, H. Melakeberhan, T.L. Niblack, W.C. Stienstra, **G.L. Tylka**, T.A. Wheeler, and D.S. Wysong. 2006. Assessing *Heterodera glycines*-resistant and susceptible cultivar yield response. *Journal of Nematology* 38:76-82.
- Tabor, G.M., S.R. Cianzio, **G.L. Tylka**, R. Roorda, and C.R. Bronson. 2006. A new greenhouse method to assay soybean resistance to brown stem rot. *Plant Disease*: 90:1186-1194.
- Leon, R., M.D.K. Owen, D. Soh, and **G.L. Tylka**. 2005. Absence of interactive responses of early soybean growth to soybean cyst nematode, post-emergence herbicides, and soil pH and texture. *Weed Technology* 19:847-854.
- Charlson, D.V. and **G.L. Tylka**. 2003. *Heterodera glycines* cyst components and surface disinfection affect *H. glycines* hatching. *Journal of Nematology* 35:458-464.
- Tabor, G.M., **G.L. Tylka**, S.C. Cianzio, and C.R. Bronson. 2003. Resistance to *Phialophora gregata* is expressed in the stems of resistant soybeans. *Plant Disease* 87:970-976.
- Tabor, G.M., **G.L. Tylka**, J.E. Behm, and C.R. Bronson. 2003. *Heterodera glycines* infection increases incidence and severity of brown stem rot of soybeans. *Plant Disease* 87:655-661.
- Wang, J., T.L. Niblack, J.N. Tremaine, W.J. Wiebold, **G.L. Tylka**, C.C. Marett, G.R. Noel, O. Myers, and M.E. Schmidt. 2003. The soybean cyst nematode reduces soybean yield without causing obvious symptoms. *Plant Disease* 87:623-628.
- Fallick, J.B., W.D. Batchelor, **G.L. Tylka**, T.L. Niblack, and J.O. Paz. 2002. Coupling soybean cyst nematode damage to CROPGRO-Soybean. *Transactions of the American Society of Agricultural Engineers* 45:433-441.
- Niblack, T.L., P.R. Arelli, G.R. Noel, C.H. Opperman, J.H. Orf, D.P. Schmitt, J.G. Shannon and **G.L. Tylka**. 2002. A new classification scheme for genetically diverse populations of *Heterodera glycines*. *Journal of Nematology* 34:279-288.
- Nutter, F.W., **G.L. Tylka**, J. Guan, A.J.D. Moreira, C.C. Marett, T.R. Rosburg, J.P. Basart, and C.S. Chong. 2002. Use of remote sensing to detect soybean cyst nematode-induced plant stress. *Journal of Nematology* 34:222-231.
- Perry, R.N., J. Beane, C.C. Marett, and **G.L. Tylka**. 2002. Comparison of the rate of embryonic development of *Globodera rostochiensis* and *G. pallida* using flow cytometric analysis. *Nematology* 4:553-555.
- Gavassoni, W.L., **G.L. Tylka**, and G.P. Munkvold. 2001. Relationship between tillage and spatial patterns of *Heterodera glycines*. *Phytopathology* 91:534-545.

## PUBLICATIONS (continued)

- Paz, J.O., W.D. Batchelor, **G.L. Tylka**, and R.G. Hartzler. 2001. A modeling approach to quantify the effects of spatial soybean yield limiting factors. *Transactions of the American Society of Agricultural Engineers* 44:1329-1334.
- Paz, J.O., W.D. Batchelor, and **G.L. Tylka**. 2001. Method to use crop growth models to estimate potential return for variable-rate management in soybeans. *Transactions of the American Society of Agricultural Engineers* 44:1335-1341.
- Buckelew, L.D., L.P. Pedigo, H.M. Mero, M.D.K. Owen, and **G.L. Tylka**. 2000. Effects of weed management systems on canopy insects in herbicide-resistant soybeans. *Journal of Economic Entomology* 93:1437-1443.
- Wang, J., P.A. Donald, T.L. Niblack, G.W. Bird, J. Faghihi, J.M. Ferris, D.J. Jardine, C. Grau, P.E. Lipps, A.E. MacGuidwin, H. Melakeberhan, G.R. Noel, P. Pierson, R.M. Reidel, P.R. Sellers, W.C. Stienstra, T.C. Todd, **G.L. Tylka**, and D.S. Wysong. 2000. Soybean cyst nematode reproduction in the north central United States. *Plant Disease* 84:77-82.
- Workneh, F., X.B. Yang, and **G.L. Tylka**. 1999. Soybean brown stem rot, *Phytophthora sojae*, and *Heterodera glycines* affected by soil texture and tillage relations. *Phytopathology* 89:844-850.
- Workneh, F., **G.L. Tylka**, X.B. Yang, J. Faghihi, and J.M. Ferris. 1999. Regional assessment of soybean brown stem rot, *Phytophthora sojae*, and *Heterodera glycines* using area-frame sampling: prevalence and effects of tillage. *Phytopathology* 89:204-211.
- Workneh, F., X.B. Yang, and **G.L. Tylka**. 1998. Effect of tillage practices on vertical distribution of *Phytophthora sojae*. *Plant Disease* 82:1258-1263.
- Levene, B.C., M.D.K. Owen, and **G.L. Tylka**. 1998. Influence of herbicide application to soybean on soybean cyst nematode egg hatching. *Journal of Nematology* 30:347-352.
- Levene, B.C., M.D.K. Owen, and **G.L. Tylka**. 1998. Response of soybean cyst nematodes and soybeans (*Glycine max*) to herbicides. *Weed Science* 46:264-270.
- Thompson, J.M., and **G.L. Tylka**. 1997. Differences in hatching of *Heterodera glycines* egg-mass and encysted eggs in vitro. *Journal of Nematology* 29:315-321.
- Kraus, G.A., S. Vander Louw, **G.L. Tylka**, and D.H. Soh. 1996. The synthesis and testing of compounds that inhibit soybean cyst nematode egg hatch. *Journal of Agricultural and Food Chemistry* 44:1548-1550.
- Shapiro, D.I., **G.L. Tylka**, and L.C. Lewis. 1996. Effects of fertilizers on virulence of *Steinernema carpocapsae*. *Applied Soil Ecology* 3:27-34.

## PUBLICATIONS (continued)

- Behm, J.E., **G.L. Tylka**, T.L. Niblack, W.J. Wiebold, and P.A. Donald. 1995. Effects of zinc fertilization of corn on hatching of *Heterodera glycines* in soil. *Journal of Nematology* 27:164-171.
- Shapiro, D.I., **G.L. Tylka**, E.C. Berry, and L.C. Lewis. 1995. Effects of earthworms on the dispersal of *Steinernema* spp. *Journal of Nematology* 27:21-28.
- Browde, J.A., L.P. Pedigo, M.D.K. Owen, **G.L. Tylka**, and B.C. Levene. 1994. Growth of soybean stressed by nematodes, herbicides, and simulated insect defoliation. *Agronomy Journal* 86:968-974.
- Browde, J.A., L.P. Pedigo, M.D.K. Owen, and **G.L. Tylka**. 1994. Soybean yield and pest management as influenced by nematodes, herbicides, and defoliating insects. *Agronomy Journal* 86:601-608.
- Browde, J.A., **G.L. Tylka**, L.P. Pedigo, and M.D.K. Owen. 1994. A method for infesting small field plots with *Heterodera glycines*. *Agronomy Journal* 86:585-587.
- Browde, J.A., **G.L. Tylka**, L.P. Pedigo, and M.D.K. Owen. 1994. Responses of *Heterodera glycines* populations to a postemergence herbicide mix and simulated insect defoliation. *Journal of Nematology* 26:498-504.
- Kraus, G.A., B. Johnston, A. Kongsjahju, and **G.L. Tylka**. 1994. The synthesis and evaluation of compounds that affect soybean cyst nematode egg hatch. *Journal of Agricultural and Food Chemistry* 42:1839-1840.
- Wong, A.T.S. and **G.L. Tylka**. 1994. Eight nonhost weed species of *Heterodera glycines* in Iowa. *Plant Disease* 78:365-367.
- Tylka, G.L.**, T.L. Niblack, T.C. Walk, K.R. Harkins, L. Barnett, and N.K. Baker. 1993. Flow cytometric analysis and sorting of *Heterodera glycines* eggs. *Journal of Nematology* 25:596-602.
- Wong, A.T., **G.L. Tylka**, and R.G. Hartzler. 1993. Effects of eight herbicides on in vitro hatching of *Heterodera glycines*. *Journal of Nematology* 25:578-584.
- Tylka, G.L.**, R.S. Hussey, and R.W. Roncadori. 1991. Axenic germination of vesicular-arbuscular mycorrhizal fungi: effects of selected *Streptomyces* species. *Phytopathology* 81:754-759.
- Tylka, G.L.**, R.S. Hussey, and R.W. Roncadori. 1991. Interactions of vesicular-arbuscular mycorrhizal fungi, phosphorus, and *Heterodera glycines* on soybean. *Journal of Nematology* 23:122-133.



## PUBLICATIONS (continued)

## II. Non-refereed, online research articles:

**Tylka, G.L.**, G.D. Gebhart, and C.C. Marett. 2006. Iowa 2005 soybean cyst nematode-resistant soybean variety trial results. *Crop Management* (<http://www.plantmanagementnetwork.org/pub/cm/trials/2005/soy/Tylka.xls>).

**Tylka, G.L.**, G.D. Gebhart, and C.C. Marett. 2006. Iowa 2004 soybean cyst nematode-resistant soybean variety trial results. *Crop Management* (<http://www.plantmanagementnetwork.org/pub/cm/trials/2004/soy/Tylka.xls>).

**Tylka, G.L.**, G.D. Gebhart, and C.C. Marett. 2006. Iowa 2003 soybean cyst nematode-resistant soybean variety trial results. *Crop Management* (<http://www.plantmanagementnetwork.org/pub/cm/trials/2003/soy/Tylka.xls>).

**Tylka, G.L.**, G.D. Gebhart, and C.C. Marett. 2003. Iowa 2002 soybean cyst nematode-resistant soybean variety trial results. *Crop Management* (<http://www.plantmanagementnetwork.org/pub/cm/trials/2002/soy/Tylka.xls>).

**Tylka, G.L.**, G.D. Gebhart, and C.C. Marett. 2003. Iowa 2001 soybean cyst nematode-resistant soybean variety trial results. *Crop Management* (<http://www.plantmanagementnetwork.org/pub/cm/trials/2001/soy/Tylka.xls>).

## III. Miscellaneous refereed online journal articles:

Robertson, A. and **G.L. Tylka**. 2007. Building the “Rust Fast Track System” for identifying Asian soybean rust in Iowa. *Journal of Extension* 45(3) Article 3IAW7. [www.joe.org/joe/2007june/iw7.shtml](http://www.joe.org/joe/2007june/iw7.shtml).

**Tylka, G.L.** and C.A. Jasalavich. 2001. Free-living and plant-parasitic nematodes (roundworms). *The Plant Health Instructor*. DOI: 10.1094/PHI-K-2001-0409-01.

Davis, E.L. and **G.L. Tylka**. 2000. Soybean cyst nematode disease. *The Plant Health Instructor*. DOI: 10.1094/PHI-I-2000-0725-01.

## IV. Book chapters:

Niblack, T.L., K.N. Lambert, and **G.L. Tylka**. 2006. A model plant pathogen from the kingdom Animalia: *Heterodera glycines*, the soybean cyst nematode. *Annual Review of Phytopathology* 44:283-303.

Kraus, G.A., **G.L. Tylka**, S. Van der Louw, and P.K. Choudhury. 2004. Chapter 11, Management of the soybean cyst nematode by using a biorational strategy, pp. 161-172 in *Agricultural Applications in Green Chemistry*, W.M. Nelson, ed., American Chemical Society, Washington, DC.

## PUBLICATIONS (continued)

- Tylka, G.L.** 2004. Management of nematode diseases: options in Encyclopedia of Plant and Crop Science. R.M. Goodman, ed., Marcel Dekker, Inc., New York.
- Niblack, T.L., **G.L. Tylka**, and R.D. Riggs. 2004. Chapter 16, Nematode pathogens of soybean, pp. 821-851 in Soybean Monograph, 3<sup>rd</sup> Edition. H.R. Boerma & J.E. Specht, eds., American Society of Agronomy, Inc., Crop Science Society of America, Inc., Soil Science Society of America, Inc., Madison, WI.
- Jones, P.W., **G.L. Tylka**, and R.N. Perry. 1998. Chapter 8, Hatching, pp. 181-212 in Physiology and Biochemistry of Free-living and Plant-parasitic Nematodes. R.N. Perry and D.J. Wright, eds., CAB International, Oxon, United Kingdom.
- V. Research abstracts (from last 5 years, 2009-present):
- Olson, L.K., **G.L. Tylka**, J. Jordhal, S. Meyer, J. Goltz, J. Kringler, T. Helms, and S. Markell. 2013. Increasing awareness of soybean cyst nematode in North Dakota. *Phytopathology* 103(S4):107-108.
- Tatalovic, N., **G.L. Tylka**, and L.F. Leandro. 2013. Effect of watering regime and *Fusarium virguliforme* (Fv) infection on location of soybean cyst nematode (SCN) syncytia in soybean roots. *Phytopathology* 103(S4):143.
- Tylka, G.L.**, M.T. McCarville, C.C. Marett, G.D. Gebhart, D.H. Soh, M.P. Mullaney, and M.E. O'Neal. 2013. Direct comparison of soybean cyst nematode reproduction on resistant soybean varieties in greenhouse and field experiments. *Journal of Nematology* 45(4): 279–330.
- Diaz-Arias, M., **G.L. Tylka**, L. Leandro, and G.P. Munkvold. 2011. Interactions between *Fusarium* root rot pathogens and *Heterodera glycines*, on soybean roots. *Phytopathology* 101:S42.
- Niblack, T.L., J. Bond, and **G.L. Tylka**. 2011. The nightmare of plant diseases associated with soybean cyst nematodes. *Phytopathology* 101:S209.
- Tatalovic, N., **G.L. Tylka**, and L.F. Leandro. 2011. Microscopic observation of the interaction between the soybean sudden death syndrome pathogen, *Fusarium virguliforme*, and the soybean cyst nematode, *Heterodera glycines*, in soybean roots. *Phytopathology* 101:S175.
- Wiggs, S.N. and **G.L. Tylka**. 2011. The nature of the relationship between soybean cyst nematode population densities and soil pH. *Phytopathology* 101:S191.
- Arias, M.M.D., **G. Tylka**, L. Leandro, and G. Munkvold. 2010. Effects of soybean cyst nematode infestation and resistance on *Fusarium* root rot on soybeans. *Phytopathology* 100:S30.
- Tylka, G.L.** 2010. Maximizing effectiveness of extension education efforts. *Journal of Nematology* 42:274-275.

## PUBLICATIONS (continued)

## VI. Extension publications:

**Tylka, G.L.**, G.D. Gebhart, C.C. Marett, M.P. Mullaney, and S.N. Wiggs. 2013. Evaluation of soybean varieties resistant to soybean cyst nematode in Iowa – 2012. Iowa State University Extension, publication IPM-52. (updated annually)

**Tylka, G.L.** 2012. Soybean cyst nematode field guide, 2<sup>nd</sup> Edition. Iowa State University Extension Publication CSI 0012, 62 pp.

**Tylka, G.L.** and M.P. Mullaney. 2012. Soybean cyst nematode-resistant soybeans for Iowa. Iowa State University Extension Publication PM 1649, 27 pp. (updated annually)

Mueller, D., A. Sisson, E. Hodgson, A. Mallarino, C. McGrath, M. O’Neal, P. Pedersen, C. Pilcher, R. Pope, M. Rice, A. Robertson, J. Sawyer, K. Schaefer, K. Simon, **G. Tylka**, and D. Wright. 2011. Soybean field guide, 2nd Edition. Iowa State University Extension Publication CSI 010, 68 pp.

Mueller, D., A. Robertson, A. Sisson, and **G. Tylka**. 2010. Soybean diseases. Iowa State University Extension Publication CSI 0004, 36 pp.

Mueller, D., R. Pope, L. Abendroth, R. Elmore, B. Hartzler, C. McGrath, G. Munkvold, M. Rice, A. Robertson, J. Sawyer, K. Schaefer, J. Tollefson, and **G. Tylka**. 2009. Corn field guide. Iowa State University Extension, publication CSI 0001, 84 pp.

Mueller, D.S., A. Robertson, and **G.L. Tylka**. 2008. Common soybean leaf diseases and Asian soybean rust. Iowa State University Extension, Publication PM 1989, 2 pp.

Mueller, D., **G. Tylka**, A. Robertson, M. O’Neal, C. Pilcher, P. Pedersen, K. Schaefer, M. Rice, A. Mallarino, R. Pope, C. McGrath, K. Simon, D. Wright, and G. Usovsky. 2008. Soybean disease and pest management field guide. Iowa State University Extension, publication CSI 0010, 52 pp.

Niblack, T.L. and **G.L. Tylka**, eds. 2008. Soybean cyst nematode management guide, 5<sup>th</sup> edition. North Central Soybean Research publication. Columbia: University of Missouri Printing Services. 18 pp.

**Tylka, G.L.** 2008. Soybean cyst nematode management field guide. Iowa State University Extension, publication CSI 0012, 56 pp.

Niblack, T.L. and **G.L. Tylka**, eds. 2003. Soybean cyst nematode management guide, 4<sup>th</sup> edition. North Central Soybean Research publication. Columbia: University of Missouri Printing Services. 18 pp.

PUBLICATIONS (continued)

Ally, M., W. Bailey, A. Blaine, C. Boerboom, M. Draper, J. Dunphy, R. Elmore, F. Fishel, C. Grau, D. Hershman, B. Johnson, S. Killpack, S. Naeve, E. Nafziger, D. Oliver, D. Peterson, G. Rehm, P. Scharf, L. Sweets, **G. Tylka**, and W. Wiebold. 2001. U.S. soybean diagnostic guide. United Soybean Board publication, 50 pp., plus CD and online at [www.psu.missouri.edu/soydoc/](http://www.psu.missouri.edu/soydoc/).

**Tylka, G.L.** and P.H. Flynn. 1999. Interpreting SCN soil sample results. Iowa State University Extension, publication IPM-61, 4 pp.

**Tylka, G.L.**, and P.E. Pierson. 1998. Getting to the root of the problem with soybean cyst nematode: the SCN Coalition grower's workbook. Iowa State University Extension, publication EDC-151, 8 pp.

**Tylka, G.L.**, and P.E. Pierson. 1998. SCN facts. Iowa State University Extension, publication EDC-150, 2 pp.

**Tylka, G.L.**, and P.E. Pierson. 1998. Scouting for SCN. Iowa State University Extension, publication EDC-149, pocket card.

**Tylka, G.L.**, and P.E. Pierson. 1998. Take the test, beat the pest: the SCN Coalition leaders guide for SCN management training. Iowa State University Extension, publication NCR-608, 35 pp. plus 104 35mm slides.

**Tylka, G.L.** 1997. Scouting for corn nematodes. Iowa State University Extension, publication IPM-53s, 1p.

**Tylka, G.L.** 1996. Scouting for soybean cyst nematode. Iowa State University Extension, publication IPM-47s, 1p.

**Tylka, G.L.** 1995. Soybean cyst nematode. Iowa State University Extension, publication Pm-879, 6 pp.

VII. Electronic extension education materials:

**Tylka, G.L.** and A. Ciha. 2009. Corn nematodes. Iowa State University Crop Adviser Institute training module (online and on CD).

**Tylka, G.L.** 2007. Soybean cyst nematode: biology, scouting, and management (presentation with audio). Online. Plant Health Management Network, Focus on Soybean. [www.plantmanagementnetwork.org/infocenter/topic/focusonsoybean/](http://www.plantmanagementnetwork.org/infocenter/topic/focusonsoybean/).

**Tylka, G.L.** and B. Brueland. 2006. Soybean cyst nematode. Iowa State University Crop Adviser Institute training module (online and on CD).

NOTEWORTHY INVITED PRESENTATIONS (last 5 years, 2009-present)

- January 2014 Syngenta National Clariva Launch Conference, January 7, 2014, Orlando, Florida, "Status of resistant soybean varieties for management of SCN"
- August 2013 Innovations in Seed Treatments for Crop Protection and Health symposium, American Phytopathological Society annual meeting, August 14, 2013, Austin, Texas, "Nematode-protectant seed treatments: New options for nematode management in row crops"
- March 2013 Soybean Cyst Nematode and Nematodes on Corn Workshop, Fargo, North Dakota, March 6-7, 2013
- February 2013 Syngenta Soybean Crop Team Nematode Conference, February 5, 2013, Campinas, Brazil, "Managing SCN with resistant soybean varieties"
- February 2013 World Soybean Research Conference, February 20, 2013, Durban, South Africa, "SCN resistance: how it all comes together in the Midwestern United States"
- March 2012 "Nematode Seed Treatment Protectants: Do Growers Need That Type of Insurance?", 7<sup>th</sup> International Integrated Pest Management Symposium, Memphis, Tennessee, March 28, 2012
- March 2012 "SCN-resistant Soybeans, HG types, Yield, and SCN Reproduction – How It All Comes Together in the Field in Iowa", Annual meeting of the Southern Soybean Disease Workers, Pensacola, Florida, March 7, 2012
- February 2012 "Soybean Cyst Nematode Update for the Region", Advanced Crop Advisers Workshop, Fargo, North Dakota, February 10, 2012
- February 2012 "Nematodes of Soybean and Corn: Recognizing the Risk and Tapping into the Right Management Tools", Advanced Crop Advisers Workshop, Fargo, North Dakota, February 9, 2012
- December 2011 "Nematodes That Feed on Corn: Prospects for 2012", South Dakota Agronomy Conference, Sioux Falls, South Dakota, December 15, 2011
- November 2011 "Nematodes That Feed on Corn: What to Make of it All", University of Missouri Crop Management Conference, Columbia, Missouri, November 30, 2011
- November 2011 "The ISU Corn and Soybean Initiative: Redefining Crops Extension Through Formal Partnerships with Private Industry Service Providers", Ohio State University, Columbus, Ohio, November 1, 2011

NOTEWORTHY INVITED PRESENTATIONS (last 5 years, 2009-present) (continued)

- November 2011      “Soybean Cyst Nematode, Host Resistance, and HG Types: What Does it All Mean for Growers?”, Ohio State University, Columbus, Ohio, November 1, 2011
- March 2011        Soybean Cyst Nematode and Nematodes on Corn Workshop, Fargo, North Dakota, March 9-10, 2011
- February 2011     “Biology and Management of Plant-parasitic Nematodes on Crop Plants”, Great Plains Consultants Meeting, Bayer CropScience, Denver, CO, February 22, 2011
- December 2010    “Biology and Management of Nematodes that Feed on Corn” and “Soybean Cyst Nematode; Biology, Scouting, and Management for South Dakota”, 2010 South Dakota Agronomy Conference, Sioux Falls, SD, December 14, 2010
- July 2010         “Maximizing Effectiveness of Extension Education Efforts”, Symposium on Educational Strategies and Methodologies in Nematology, Society of Nematologists 2010 Annual Meeting, Boise, ID, July 13, 2010
- February 2009     “Managing Soybean Cyst Nematode Using Precision Farming Technologies”, National Alliance of Independent Crop Consultants, Bloomington, MN, February 12, 2009

PATENTS AWARDED

Coats, J. R., A.L. Egger, C.J. Peterson, R. Tsao, and **G.L. Tylka**. 2003. Compounds related to natural sources and their use as biopesticides. U.S. Patent Number 6,545,043.

Coats, J. R., C.J. Peterson, R. Tsao, A.L. Egger, and **G.L. Tylka**. 2001. Biopesticides related to natural sources. U.S. Patent Number 6,207,705.

Kraus, G.A., **G.L. Tylka**, and S. Van der Louw. 1997. Ketodiacid compounds that inhibit nematode egg hatching. U.S. Patent Number 5,648,318.

BEFORE THE PUBLIC UTILITIES COMMISSION  
OF THE STATE OF SOUTH DAKOTA

IN THE MATTER OF THE APPLICATION OF  
MONTANA-DAKOTA UTILITIES CO. AND  
OTTER TAIL POWER COMPANY FOR A  
PERMIT TO CONSTRUCT THE BIG STONE  
TO SOUTH ELLENDALE 345KV  
TRANSMISSION LINE

AMENDED CERTIFICATE OF SERVICE FOR  
DIRECT TESTIMONY OF  
GREGORY TYLKA

EL13-028

**CERTIFICATE OF SERVICE**

The undersigned attorney hereby certifies that a true and correct copy of the **DIRECT TESTIMONY OF GREGORY TYLKA** was served upon the following parties of record, electronically or in paper form, this 24<sup>th</sup> day of April, 2014:

Thomas J. Welk, Attorney  
P.O. Box 5015  
Sioux Falls, SD 57117-5015  
[tjwelk@bgpw.com](mailto:tjwelk@bgpw.com)

Darren Kearny  
Staff Analyst  
South Dakota P.U.C.  
500 E. Capitol Ave.  
Pierre, SD 57501  
[darren.kearny@state.sd.us](mailto:darren.kearny@state.sd.us)

Sandra Raap  
Day County Auditor  
711 W. First St., Ste. 204  
Webster, SD 57274  
[dcaud@itctel.com](mailto:dcaud@itctel.com)

Patricia Van Gerpen  
Executive Director  
South Dakota P.U.C.  
500 E. Capitol Ave.  
Pierre, SD 57501  
[patty.vangerpen@state.sd.us](mailto:patty.vangerpen@state.sd.us)

Jennifer Smestad, Attorney  
214 S. Cascade St.  
Fergus Falls, MN 56538  
[jsmestad@ottertail.com](mailto:jsmestad@ottertail.com)

Karen Layher  
Grant County Auditor  
210 E. Fifth Ave.  
Milbank, SD 57252  
[karen.layher@state.sd.us](mailto:karen.layher@state.sd.us)


Karen Cremer, Attorney  
500 E. Capitol Ave.  
Pierre, SD 57501  
[karen.cremer@state.sd.us](mailto:karen.cremer@state.sd.us)

Daniel S. Kuntz, Attorney  
P.O. Box 5650  
1200 West Century Ave.  
Bismarck, ND 58506  
[dan.kuntz@mduresources.com](mailto:dan.kuntz@mduresources.com)

Brian Rounds  
Staff Analyst  
South Dakota P.U.C.  
500 E. Capitol Ave.  
Pierre, SD 57501  
[brian.rounds@state.sd.us](mailto:brian.rounds@state.sd.us)

Maxine Fischer  
Brown County Auditor  
25 Market St., Ste. 1  
Aberdeen, SD 57401  
[maxine.fischer@browncounty.sd.gov](mailto:maxine.fischer@browncounty.sd.gov)

Dated this 7<sup>th</sup> Day of June, 2014



---

N. Bob Pesall, Attorney  
P.O. Box 23  
Flandreau, SD 57028  
(605) 573-0274