Federal and State Listed Threatened and Endangered Species in South Dakota

Common Name	Scientific Name	Federal Status	Federal County Listing	Potential Impact	Habitat Requirement	Determination of Effect
Mammals		4				
Northern long-eared bat	Myotis septentrionalis	PE	Beadle, Campbell, Edmunds, Faulk, Kingsbury, Lake, Lincoln, McCook, McPherson, Miner, Minnehaha, Spink, Turner	No effect	Summer roosting habitat underneath bark or in crevices of live and dead trees. Winter habitat includes caves and mines with large entrances.	The range of this species is located at the eastern border of the state (NatureServe, 2014). Forested areas within this part of the Project are limited to the Big Sioux River crossing. The Big Sioux River will be crossed via HDD, therefore no impacts to forested areas are anticipated.
Birds			1.1.1.1.1.1.1.1.1.1.1.1.1.1.1.1.1.1.1.			
Interior least tern	Sterna antillarum athalassos	E	Campbell	No effect	Interior least tern nesting habitat includes open shorelines, riverine sandbars, and mudflats along Missouri and Mississippi Rivers drainages.	The Project does not cross the Missouri River within South Dakota. No suitable habitat within the Project area.
Piping plover	Charadrius melodus	т	Campbell, Kingsbury	No effect	Sandy or gravelly beaches and sandbars or alkaline wetlands.	No suitable nesting habitat was identified during Project field surveys. Critical habitat for the piping plover is along the Missouri River; the Project doe not cross the Missouri River within South Dakota. This species is highly mobile and would likely avoid the construction area.
Red knot	Calidris canutus rufa	PT	Beadle, Campbell, Edmunds, Faulk, Kingsbury, Lake, Lincoln, McCook, McPherson, Miner, Minnehaha, Spink, Turner	No effect	Breeds in the Arctic tundra areas, such as sparsely vegetated habitat. When non- breeding they prefer primarily intertidal, marine habitats, coastal inlets, estuaries, and bays.	No suitable habitat within the Project area
Sprague's pipit	Anthus spragueii	с	Campbell, McPherson	No effect	Prefer native grasslands of intermediate height and sparse to intermediate vegetation density, low forb density, and little bare ground but low litter depth. Introduced grasslands may be utilized, but to a much lesser extent. Nests on the ground from early May to mid-October.	Breeding habitat range is in the northern part of the state. Some of the of the Project area may be within this range; however, there are no occurrences documented within the Project area (SDNHP, 2014 and eBird, 2014)

and the restart the second second

Common Name	Scientific Name	Federal Status	Federal County Listing	Potential Impact	Habitat Requirement	Determination of Effect
Whooping crane	Grus americana	E	Beadle, Campbell, Clark, Edmunds, Faulk, Kingsbury, McCook, McPherson, Miner, Spink, Turner	No effect	During migration, this species utilizes wetlands and cropland ponds for feeding and roosting. Seasonal and semi-permanent wetlands are the most commonly used.	The Project area is within the migratory range of this species (Cornell Lab of Ornithology, 2014). Only one whooping crane occurrence record is located in Kingsbury County within one mile of the Project (SDNHP, 2014). This species is highly mobile and would likely avoid construction.
Fishes						
Pallid sturgeon	Scaphirhynchus albus	E	Campbell, Lincoln	No effect	Prefer a fast flowing turbid river with a firm sand or gravel bottom. Areas at the end of chutes or sandbars are commonly used for feeding.	The Missouri River (Campbell County) will not be crossed in South Dakota, and the Big Sioux River (Lincoln County) will be crossed via HDD, therefore no impacts will occur to this species.
Topeka shiner	Notropis topeka	E	Beadle, Kingsbury, Lake, Lincoln, McCook, Miner, Minnehaha, Spink, Turner	Not likely to adversely affect	Found in small prairie streams that exhibit perennial or nearly perennial flow. Substrate usually is clean gravel, cobble, or sand.	Nine streams crossed by the Project may contain this species as identified by the USFWS. Two of these streams (James and Big Sioux Rivers) will be crossed via HDD. Consultation with the U.S. Fish an Wildlife Service is necessary. Mitigation measures to be used during construction of the pipeline within identified streams are under development.
Invertebrates		(1)世代][1]		a the second	1. 新加速 医血病	
Dakota skipper	Hesperia dacotae	Т	Edmunds, McPherson	No effect	Dakota skippers only utilize high quality undisturbed (i.e., remnant, uncultivated) prairie; including, wet tallgrass prairie and dry mixed grass prairie.	No native grasslands were identified within Edmunds and McPherson County during field surveys.
Vascular Plant	s - 110	in the second		A Dama Contract		
Western prairie fringed orchid	Platanthera praeclara	Т	Lake, Lincoln, McCook, Miner, Minnehaha, Turner	No effect	Prefers moist tallgrass prairie and sedge meadows are appropriate habitat for the western prairie fringed orchid.	Suitable habitat present within Project area, however the species seems to have been extirpated from South Dakota.
E= Endangered T= Threatened PE=Proposed E PT= Proposed T C= Candidate	ndangered					

State Listed Threatened and Endangered Species in South Dakota						
Common Name	Scientific Name	State Status <sup>a</sup>	Potential Impact	Habitat Requirement	Determination of Effect	
Mammals	HERE WALL STORES	11.20	無意か		15、2000年間電信であった。 三人名(日本)	
Black-footed ferret	Mustela nigripes	E	No impact anticipated	Associated exclusively with large (10,000 acres or more) prairie dog towns. Use burrows for shelter and feed on prairie dogs and other species within the habitat.	Historically, the species was present within the state; however, large prairie dog complexes needed to support a black-footed ferret population do not currently exist within the Project area.	
Northern river otter	Lontra canadensis	Т	No impact anticipated	Rivers with high quality water and an abundant food supply.	Within the Project area, this species has been documented within the Big Sioux River and James River watersheds (South Dakota Game, Fish, and Parks [SDGFP], 2014a and South Dakota Natural Heritage Program [SDNHP], 2014). However, both of these rivers will be crossed via HDD, therefore avoiding impacts to the riverine habitats utilized by the otter.	
Swift fox	Vulpes velox	Т	No impact anticipated	Prefer short or mixed grass prairies with flat to gently rolling terrain and sparse vegetation that allows for good mobility and visibility.	Although historically the range of this species was within the Project area, the species does not currently reside within the Project area (NatureServe, 2014).	
Birds						
American dipper	Cinclus mexicanus	Т	No impact anticipated	Cold and clear, fast-moving streams with gravel, stone, or sand bottoms which support invertebrates. Streams with structures over the water such as waterfalls, rocks and boulders are needed for nesting.	The range of this species is not within the Project area (Cornell Lab of Ornithology, 2014).	
Bald eagle	Haliaeetus leucocephalus	т	No impact anticipated	Breeds and winters in areas close to a coast, river or lake. Prefers conifers for nesting and roosting and tends to avoid areas with high human traffic.	There are few forested areas along the pipeline route for nesting. Occurrence data from the SDNHP documented a nest approximately one mile from the Project area. Field surveys did not identify bald eagles within the Project area. This species is highly mobile and would likely avoid construction.	
Eskimo curlew	Numenius borealis	E	No impact anticipated	Variety of grassland habitats.	The Project area is within the migratory range of this species (NatureServe, 2014). This species is highly mobile and would likely avoid construction.	
Interior least tern	Sterna antillarum athalassos	E	No impact anticipated	Interior least tern nesting habitat includes open shorelines, riverine sandbars, and mudflats along Missouri and Mississippi Rivers drainages.	The Project does not cross the Missouri River within South Dakota. No suitable habitat within the Project area.	
Osprey	Pandion haliaetus	т	No impact anticipated	Prefer habitat near water including, saltmarshes, rivers, ponds, and reservoirs. Osprey places their nest in open areas on poles, channel markers, and dead trees, often over water.	The Project area is within the migratory range of this species (Cornell Lab of Ornithology, 2014). This species is highly mobile and would likely avoid construction.	

Common		State	Potential		
Name	Scientific Name	Status <sup>a</sup>	Impact	Habitat Requirement	Determination of Effect
Peregrine falcon	Falco peregrines	E	No impact anticipated	Inhabits any open habitat with a wide view of the surrounding area, close proximity to water and rocky cliffs or even tall buildings available for nesting.	No nesting habitat is within the Project area (NatureServe, 2014). This species is highly mobile and would likely avoid the construction area.
Piping plover	Charadrius melodus	т	No impact anticipated	Sandy or gravelly beaches and sandbars or alkaline wetlands.	No suitable nesting habitat was identified during Project field surveys. Critical habitat for the piping plover is along the Missouri River; the Project does not cross the Missouri River within South Dakota. This species is highly mobile and would likely avoid the construction area.
Whooping crane	Grus americana	E	No impact anticipated	During migration, this species utilizes wetlands and cropland ponds for feeding and roosting. Seasonal and semi-permanent wetlands are the most commonly used.	The Project area is within the migratory range of this species (Cornell Lab of Ornithology, 2014). Only one whooping crane occurrence record is located in Kingsbury County within one mile of the Project (SDNHP, 2014). This species is highly mobile and would likely avoid construction.
Reptiles	- 193 - H	the first section of the			
Eastern hognose snake	Heterodon platirhinos	Т	No impact anticipated	Prefer woodlands with sandy soil, fields, farmland and coastal areas.	The range of this species is not located within the Project area (NatureServe, 2014).
False map turtle	Graptemys pseudogeograph ica	Т	No impact anticipated	Inhabits slow moving rivers, river sloughs, oxbow lakes, lakes and reservoirs containing abundant aquatic vegetation and basking sites.	The range of this species within South Dakota is limite to the Missouri River area. The Project enters South Dakota east of the Missouri River (NatureServe, 2014)
Lined snake	Tropidoclonion lineatum	E	No impact anticipated	Prefers open prairie hillsides and rocky, woodland areas	The range of this species within South Dakota is limite to the southeast corner of the state. Suitable habitat ma be present within the Project area; however, this specie is highly mobile and would likely avoid construction.
Fishes	: 為智聞	<b>这个明白</b> 的人名			
Banded killifish	Fundulus diaphanous	E	No impact anticipated	Habitat ranges from quiet waters of lakes and ponds with ample vegetation to muddy streams without vegetation.	The current species habitat range is not located within the Project area (SDGFP, 2014b).
Blacknose shiner	Notropis heterolepis	E	No impact anticipated	Prefers clear, cool streams with sand and gravel beds, and deep pools with abundant vegetation both in the water and on lands bordering the streams. This species has only been found in two pristine streams located in south-central South Dakota.	No suitable habitat within the Project area.

Common Name	Scientific Name	State Status <sup>a</sup>	Potential Impact	Habitat Requirement	Determination of Effect
Finescale dace	Chrosomus neogaeus	E	No impact anticipated	Occur most often in cool, clear mountain streams and less often in lakes, reservoirs, or large rivers. Prefer moderate water velocities, associate with a variety of substrates.	The Project area is outside of the current species range (NatureServe, 2014).
Longnose sucker	Catostomus catostomus	Т	No impact anticipated	Found in cool, spring-fed streams where it feeds on the bottom on crustaceans, snails, insect larvae, and larvae.	The Project area is outside of the current species range (NatureServe, 2014).
Northern pearl dace	Margariscus nachtriebi	Т	No impact anticipated	Occurs in cool bogs, ponds, lakes, and clear streams.	The species distribution is not located within the Project area. Limited to Counties within southwestern South Dakota (U.S. Geological Survey, 2014)
Northern redbelly dace	Chrosomus eos	Т	No impact anticipated	Prefers areas with beds of aquatic vegetation in spring-fed streams.	Believed to be extirpated from the Big Sioux drainage (SDGFP, 2014c)
Pallid sturgeon	Scaphirhynchus albus	E	No impact anticipated	Prefer a fast flowing turbid river with a firm sand or gravel bottom. Areas at the end of chutes or sandbars are commonly used for feeding.	The Missouri River (Campbell County) will not be crossed in South Dakota, and the Big Sioux River (Lincoln County) will be crossed via HDD, therefore no impacts will occur to this species.
Sicklefin chub	Macrhybopsis meeki	E	No impact anticipated	Prefer large, turbid rivers with a diversity of depths and velocities forming braided channels, sand bars, sand flats, and gravel bars.	No suitable habitat within the Project area.
Sturgeon chub	Macrhybopsis gelida	Т	No impact anticipated	Prefer large, turbid rivers with a range of depths and velocities forming braided channels, gravel bars, and sand flats and bars.	No suitable habitat within the Project area.

<sup>a</sup> South Dakota state listed species do not have county listings, they are listed state-wide.

## **REFERENCES:**

- Cornell Lab of Ornithology. 2014. All About Birds. http://www.allaboutbirds.org/guide/search. Accessed November and December 2014.
- eBird. 2014. eBird Observation Maps. http://ebird.org/ebird/map/. Accessed November and December 2014.

NatureServe. 2014. NatureServe Explorer. http://explorer.natureserve.org/. Accessed December 2014.

- South Dakota Natural Heritage Program. 2014. South Dakota Game, Fish, and Parks, Wildlife Diversity Program. Occurrence data within 2 miles of pipeline route. Received July 2014.
- South Dakota Game, Fish, and Parks. 2014a. Northern River Otter. http://gfp.sd.gov/wildlife/management/diversity/river-otter.aspx. Accessed December 2014.
- South Dakota Game, Fish, and Parks. 2014b. Banded Killifish. http://gfp.sd.gov/wildlife/critters/fish/rare-fish/banded-killifish.aspx. Accessed December 2014.
- South Dakota Game, Fish, and Parks. 2014c. Northern Redbelly Dace. http://gfp.sd.gov/wildlife/critters/fish/rare-fish/northern-redbelly-dace.aspx. Accessed November 2014.
- U.S. Geological Survey. 2014. Endangered, Threatened, & Rare Animals in South Dakota. http://www.npwrc.usgs.gov/resource/wildlife/sdrare/species/semomarg.htm. Accessed November 2014.